

EXHIBIT B

BEST AVAILABLE COPY



Entrez PubMed
Search PubMed

Nucleotide Protein Genome Structure OMIM PMC Journals Br
Go Clear
Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Display Abstract Show: 20 Sort Send to Text

☐ 1: J Pain Symptom Manage. 2004 Mar;27(3):196-205.

Related Articles, List



A comparison of five pain assessment scales for nursing home residents with varying degrees of cognitive impairment.

Closs SJ, Barr B, Briggs M, Cash K, Seers K.

School of Healthcare Studies, Baines Wing, University of Leeds, Leeds LS2 9UT, United Kingdom.

The aim of the study was to compare five different pain assessment scales for use with people with different levels of cognitive impairment who resided in nursing homes. The verbal rating scale, horizontal numeric rating scale, Face pictorial scale, color analogue scale and mechanical visual analogue scale were presented in random order to 113 residents. Cognitive impairment was assessed using the Mini-Mental State Examination. The use of the verbal rating scale was the most successful with this group, completed by 80.5% overall, and 36% of those with severe cognitive impairment. Repeated explanation improved completion rates for all the scales. Consistency between scores on the five scales was good for those with none to moderate cognitive impairment and poor for those severely impaired. This study showed no difference in pain scores according to cognitive status.


PMID: 15010098 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Privacy Statement | Freedom of Information Act | Disclaimer

Jul 27 2004 06:47

Handwritten signature



Entrez PubMed
Search PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Br

for
Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display Abstract

Show: 20 Sort

Send to Text

Text Version

Related Articles, List

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

☐ 1: J Pain Symptom Manage. 2001 Jun;21(6):450-5.

ELSEVIER SCIENCE
FULL-TEXT ARTICLE

Utilizing pain assessment scales increases the frequency of diagnosing pain among elderly nursing home residents.

Kamel HK, Phlavan M, Malekgoudarzi B, Gogel P, Morley JE.

Division of Geriatric Medicine, Saint Louis University School of Medicine, St. Louis, MO 63104, USA.

To study the effect of utilizing a combination of three easily-administered pain assessment instruments on the frequency of diagnosing pain among elderly nursing home residents, a cross-sectional study was conducted at two academic skilled nursing facilities: an 899-bed facility in Long Island, New York, and a 201-bed facility in Maryland Heights, Missouri. A total of 305 randomly selected elderly (≥ 60 years) subjects were enrolled in this study. The study involved medical records abstracting and patient interviews. Pain was assessed in 150 (49%) subjects using the question "Do you have pain?" (Group 1). In the remaining 155 (51%) subjects (Group 2), pain was assessed using three pain assessment instruments: the visual analog scale, the behavior (faces) scale, and the pain descriptive scale. The overall frequency of diagnosing pain was greater in the subjects in Group 2 compared to the subjects in Group 1 (30% vs. 15%, $P < 0.01$). Using the three pain assessment scales greatly increased the frequency of diagnosing pain among the old-old (≥ 85 years) residents (32% Group 2 vs. 6% Group 1, $P < 0.001$). The frequency of diagnosing pain among cognitively impaired residents showed a similar trend (16% Group 2 vs. 10% Group 1, $P = 0.2$). These data indicate that using three easily-administered pain assessment scales increased the frequency of diagnosing pain among nursing home residents.

PMID: 11397602 [PubMed - indexed for MEDLINE]

Display Abstract

Show: 20 Sort

Send to Text

Write to the Help Desk
NCBI | NLM | NIH

Department of Health & Human Services
Privacy Statement | Freedom of Information Act | Disclaimer

Handwritten signature

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.